
# **EE464: Algebra and Duality**

# **Example**

minimize 
$$x_1x_2$$
 subject to  $x_1 \ge 0$   $x_2 \ge 0$   $x_1^2 + x_2^2 \le 1$ 

- The objective is not convex.
- The Lagrange dual function is




$$g(\lambda) = \inf_{x} \left( x_1 x_2 - \lambda_1 x_1 - \lambda_2 x_2 + \lambda_3 (x_1^2 + x_2^2 - 1) \right)$$

$$= \begin{cases} -\lambda_3 - \frac{1}{2} \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix}^T \begin{bmatrix} 2\lambda_3 & 1 \\ 1 & 2\lambda_3 \end{bmatrix}^{-1} \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} & \text{if } \lambda_3 > \frac{1}{2} \\ -\infty & \text{otherwise, except bdry} \end{cases}$$

# Example, continued

The dual problem is

maximize 
$$g(\lambda)$$
 subject to  $\lambda_1 \geq 0$   $\lambda_2 \geq 0$   $\lambda_3 \geq \frac{1}{2}$ 



- By symmetry, if the maximum  $g(\lambda)$  is attained, then  $\lambda_1=\lambda_2$  at optimality
- The optimal  $g(\lambda^\star) = -\frac{1}{2}$  at  $\lambda^\star = (0,0,\frac{1}{2})$
- Here we see an example of a duality gap; the primal optimal is strictly greater than the dual optimal

## **Example, continued**

It turns out that, using the Schur complement, the dual problem may be written as

maximize 
$$\gamma$$
 subject to 
$$\begin{bmatrix} -2\gamma-2\lambda_3 & \lambda_1 & \lambda_2 \\ \lambda_1 & 2\lambda_3 & 1 \\ \lambda_2 & 1 & 2\lambda_3 \end{bmatrix} > 0$$
 
$$\lambda_1 > 0$$
 
$$\lambda_2 > 0$$

We'll see a systematic way to convert a dual problem to an SDP, whenever the objective and constraint functions are polynomials.

#### The Dual is Not Intrinsic

- The dual problem, and its corresponding optimal value, are *not properties of the primal feasible set and objective function alone*.
- Instead, they depend on the particular equations and inequalities used

To construct equivalent primal optimization problems with different duals:


- replace the objective  $f_0(x)$  by  $h(f_0(x))$  where h is increasing
- introduce new variables and associated constraints, e.g.

add redundant constraints

## **Example**

Adding the redundant constraint  $x_1x_2 \ge 0$  to the previous example gives

minimize 
$$x_1x_2$$
 subject to  $x_1 \ge 0$   $x_2 \ge 0$   $x_1^2 + x_2^2 \le 1$   $x_1x_2 \ge 0$ 



Clearly, this has the same primal feasible set and same optimal value as before

#### **Example Continued**

The Lagrange dual function is

$$\begin{split} g(\lambda) &= \inf_x \left( x_1 x_2 - \lambda_1 x_1 - \lambda_2 x_2 + \lambda_3 (x_1^2 + x_2^2 - 1) - \lambda_4 x_1 x_2 \right) \\ &= \begin{cases} -\lambda_3 - \frac{1}{2} \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix}^T \begin{bmatrix} 2\lambda_3 & 1 - \lambda_4 \\ 1 - \lambda_4 & 2\lambda_3 \end{bmatrix}^{-1} \begin{bmatrix} \lambda_1 \\ \lambda_2 \end{bmatrix} & \text{if } 2\lambda_3 > 1 - \lambda_4 \\ -\infty & \text{otherwise, except bdry} \end{cases} \end{split}$$

- Again, this problem may also be written as an SDP. The optimal value is  $g(\lambda^\star)=0$  at  $\lambda^\star=(0,0,0,1)$
- Adding redundant constraints makes the dual bound tighter
- This always happens! Such redundant constraints are called valid inequalities.

## **Constructing Valid Inequalities**

The function  $f:\mathbb{R}^n \to \mathbb{R}$  is called a *valid inequality* if

$$f(x) \ge 0$$
 for all feasible  $x$ 

Given a set of inequality constraints, we can generate others as follows.

- (i) If  $f_1$  and  $f_2$  define valid inequalities, then so does  $h(x) = f_1(x) + f_2(x)$
- (ii) If  $f_1$  and  $f_2$  define valid inequalities, then so does  $h(x) = f_1(x)f_2(x)$
- (iii) For any f, the function  $h(x) = f(x)^2$  defines a valid inequality

Now we can use *algebra* to generate valid inequalities.

## **Valid Inequalities and Cones**

- The set of *polynomial* functions on  $\mathbb{R}^n$  with real coefficients is denoted  $\mathbb{R}[x_1,\ldots,x_n]$
- Computationally, they are easy to parametrize. We will consider polynomial constraint functions.

A set of polynomials  $P \subset \mathbb{R}[x_1, \dots, x_n]$  is called a *cone* if

- (i)  $f_1 \in P$  and  $f_2 \in P$  implies  $f_1 f_2 \in P$
- (ii)  $f_1 \in P$  and  $f_2 \in P$  implies  $f_1 + f_2 \in P$
- (iii)  $f \in \mathbb{R}[x_1, \dots, x_n]$  implies  $f^2 \in P$

It is called a *proper cone* if  $-1 \notin P$ 

By applying the above rules to the inequality constraint functions, we can generate a *cone* of valid inequalities

#### **Algebraic Geometry**

- There is a correspondence between the *geometric object* (the feasible subset of  $\mathbb{R}^n$ ) and the *algebraic object* (the cone of valid inequalities)
- This is a dual relationship; we'll see more of this later.
- The dual problem is constructed from the cone.
- For equality constraints, there is another algebraic object; the *ideal* generated by the equality constraints.
- For optimization, we need to look both at the geometric objects (for the primal) and the algebraic objects (for the dual problem)

#### Cones

• For  $S \subset \mathbb{R}^n$ , the cone defined by S is

$$C(S) = \left\{ f \in \mathbb{R}[x_1, \dots, x_n] \mid f(x) \ge 0 \text{ for all } x \in S \right\}$$

- If  $P_1$  and  $P_2$  are cones, then so is  $P_1 \cap P_2$
- A polynomial  $f \in \mathbb{R}[x_1, \dots, x_n]$  is called a *sum-of-squares* (SOS) if

$$f(x) = \sum_{i=1}^{r} s_i(x)^2$$

for some polynomials  $s_1, \ldots, s_r$  and some  $r \geq 0$ . The set of SOS polynomials  $\Sigma$  is a cone.

• Every cone contains  $\Sigma$ .

#### Cones

The set  $\mathbf{monoid}\{f_1, \dots, f_m\} \subset \mathbb{R}[x_1, \dots, x_n]$  is the set of all finite products of polynomials  $f_i$ , together with 1.

The smallest cone containing the polynomials  $f_1, \ldots, f_m$  is

$$\mathbf{cone}\{f_1,\ldots,f_m\} = \left\{ \sum_{i=1}^r s_i g_i \mid s_0,\ldots,s_r \in \Sigma, \right.$$

$$g_i \in \mathbf{monoid}\{f_1, \dots, f_m\}$$

 $\mathbf{cone}\{f_1,\ldots,f_m\}$  is called the *cone generated by*  $f_1,\ldots,f_m$ 

## **Explicit Parametrization of the Cone**

- If  $f_1, \ldots, f_m$  are valid inequalities, then so is every polynomial in  $\mathbf{cone}\{f_1, \ldots, f_m\}$
- The polynomial h is an element of  $\mathbf{cone}\{f_1,\ldots,f_m\}$  if and only if

$$h(x) = s_0(x) + \sum_{i=1}^m s_i(x)f_i(x) + \sum_{i \neq j} r_{ij}(x)f_i(x)f_j(x) + \dots$$

where the  $s_i$  and  $r_{ij}$  are sums-of-squares.

## An Algebraic Approach to Duality

Suppose  $f_1, \ldots, f_m$  are polynomials, and consider the feasibility problem

does there exist  $x \in \mathbb{R}^n$  such that  $f_i(x) \ge 0$  for all  $i = 1, \dots, m$ 

Every polynomial in  $\mathbf{cone}\{f_1,\ldots,f_m\}$  is non-negative on the feasible set.

So if there is a polynomial  $q \in \mathbf{cone}\{f_1, \dots, f_m\}$  which satisfies

$$q(x) \le -\varepsilon < 0$$
 for all  $x \in \mathbb{R}^n$ 

then the primal problem is infeasible.

## **Example**

Let's look at the feasibility version of the previous problem. Given  $t \in \mathbb{R}$ , does there exist  $x \in \mathbb{R}^2$  such that

$$x_1 x_2 \le t$$

$$x_1^2 + x_2^2 \le 1$$

$$x_1 \ge 0$$

$$x_2 \ge 0$$

Equivalently, is the set S nonempty, where

$$S = \left\{ x \in \mathbb{R}^n \mid f_i(x) \ge 0 \text{ for all } i = 1, \dots, m \right\}$$

where

$$f_1(x) = t - x_1 x_2$$
  $f_2(x) = 1 - x_1^2 - x_2^2$   
 $f_3(x) = x_1$   $f_4(x) = x_2$ 

## **Example Continued**

Let  $q(x) = f_1(x) + \frac{1}{2}f_2(x)$ . Then clearly  $q \in \mathbf{cone}\{f_1, f_2, f_3, f_4\}$  and

$$q(x) = t - x_1 x_2 + \frac{1}{2} (1 - x_1^2 - x_2^2)$$

$$= t + \frac{1}{2} - \frac{1}{2} (x_1 + x_2)^2$$

$$\leq t + \frac{1}{2}$$

So for any  $t < -\frac{1}{2}$ , the primal problem is infeasible.

This corresponds to Lagrange multipliers  $(1,\frac{1}{2})$  for the thm. of alternatives.

Alternatively, this is a proof by contradiction.

- If there exists x such that  $f_i(x) \geq 0$  for  $i = 1, \ldots, 4$  then we must also have  $q(x) \geq 0$ , since  $q \in \mathbf{cone}\{f_1, \ldots, f_4\}$
- But we proved that q is negative if  $t<-\frac{1}{2}$

## **Example Continued**

We can also do better by using other functions in the cone. Try

$$q(x) = f_1(x) + f_3(x)f_4(x)$$
$$= t$$

giving the stronger result that for any t < 0 the inequalities are infeasible. Again, this corresponds to Lagrange multipliers (1,1)

- In both of these examples, we found q in the cone which was globally negative. We can view q as the Lagrangian function evaluated at a particular value of  $\lambda$
- The Lagrange multiplier procedure is *searching* over a *particular subset* of functions in the cone; those which are generated by *linear combinations* of the original constraints.
- By searching over more functions in the cone we can do better

#### **Normalization**

In the above example, we have

$$q(x) = t + \frac{1}{2} - \frac{1}{2}(x_1 + x_2)^2$$

We can also show that  $-1 \in \mathbf{cone}\{f_1, \dots, f_4\}$ , which gives a very simple proof of primal infeasibility.

Because, for  $t < -\frac{1}{2}$ , we have

$$-1 = a_0 q(x) + a_1 (x_1 + x_2)^2$$

and by construction q is in the cone, and  $(x_1+x_2)^2$  is a sum of squares.

Here  $a_0$  and  $a_1$  are positive constants

$$a_0 = \frac{-2}{2t+1} \qquad a_1 = \frac{-1}{2t+1}$$

## **An Algebraic Dual Problem**

Suppose  $f_1, \ldots, f_m$  are polynomials. The primal feasibility problem is

does there exist 
$$x \in \mathbb{R}^n$$
 such that  $f_i(x) \geq 0$  for all  $i = 1, \dots, m$ 

The dual feasibility problem is

Is it true that 
$$-1 \in \mathbf{cone}\{f_1, \dots, f_m\}$$

If the dual problem is feasible, then the primal problem is infeasible.

In fact, a result called the *Positivstellensatz* implies that *strong duality* holds here.

## **Interpretation: Searching the Cone**

Lagrange duality is searching over linear combinations with nonnegative coefficients

$$\lambda_1 f_1 + \cdots + \lambda_m f_m$$

to find a globally negative function as a certificate

• The above algebraic procedure is searching over *conic combinations* 

$$s_0(x) + \sum_{i=1}^m s_i(x)f_i(x) + \sum_{i \neq j} r_{ij}(x)f_i(x)f_j(x) + \dots$$

where the  $s_i$  and  $r_{ij}$  are sums-of-squares

## **Interpretation: Formal Proof**

We can also view this as a type of *formal proof*:

- View  $f_1, \ldots, f_m$  are *predicates*, with  $f_i(x) \ge 0$  meaning that x satisfies  $f_i$ .
- Then  $\mathbf{cone}\{f_1,\ldots,f_m\}$  consists of predicates which are *logical consequences* of  $f_1,\ldots,f_m$ .
- If we find -1 in the cone, then we have a proof by contradiction.

Our objective is to *automatically search* the cone for negative functions; i.e., proofs of infeasibility.

# **Example: Linear Inequalities**

Does there exist  $x \in \mathbb{R}^n$  such that

$$Ax \ge 0$$

$$c^T x \le -1$$

Write A in terms of its rows  $A = \begin{bmatrix} a_1^T \\ \vdots \\ a_m^T \end{bmatrix}$  ,

then we have inequality constraints defined by linear polynomials

$$f_i(x) = a_i^T x \qquad \text{for } i = 1, \dots, m$$
  
$$f_{m+1}(x) = -1 - c^T x$$

## **Example: Linear Inequalities**

We'll search over functions  $q \in \mathbf{cone}\{f_1, \dots, f_{m+1}\}$  of the form

$$q(x) = \sum_{i=1}^{m} \lambda_i f_i(x) + \mu f_{m+1}(x)$$

Then the algebraic form of the dual is:

does there exist 
$$\lambda_i \geq 0$$
,  $\mu \geq 0$  such that 
$$q(x) = -1 \qquad \text{for all } x$$

if the dual is feasible, then the primal problem is infeasible

## **Example: Linear Inequalities**

The above dual condition is

$$\lambda^T A x + \mu(-1 - c^T x) = -1 \qquad \text{for all } x$$

which holds if and only if  $A^T\lambda = \mu c$  and  $\mu = 1$ .

So we can state the duality result as follows.

#### Farkas Lemma

If there exists  $\lambda \in \mathbb{R}^m$  such that

$$A^T \lambda = c \qquad \text{and} \qquad \lambda \ge 0$$

then there does not exist  $x \in \mathbb{R}^n$  such that

$$Ax \ge 0$$
 and  $c^T x \le -1$ 

#### **Farkas Lemma**

Farkas Lemma states that the following are strong alternatives

- (i) there exists  $\lambda \in \mathbb{R}^m$  such that  $A^T \lambda = c$  and  $\lambda \geq 0$
- (ii) there exists  $x \in \mathbb{R}^n$  such that  $Ax \ge 0$  and  $c^Tx < 0$

Since this is just Lagrangian duality, there is a geometric interpretation


(i) c is in the convex cone

$$\{A^T\lambda \mid \lambda \geq 0\}$$

(ii) x defines the hyperplane

$$\{ y \in \mathbb{R}^n \mid y^T x = 0 \}$$

which separates c from the cone



## **Optimization Problems**

Let's return to optimization problems instead of feasibility problems.

minimize 
$$f_0(x)$$
 subject to  $f_i(x) \geq 0$  for all  $i=1,\ldots,m$ 

The corresponding feasibility problem is

$$t - f_0(x) \ge 0$$
 for all  $i = 1, \dots, m$ 

One simple dual is to find polynomials  $s_i$  and  $r_{ij}$  such that

$$t - f_0(x) + \sum_{i=1}^m s_i(x)f_i(x) + \sum_{i \neq j} r_{ij}(x)f_i(x)f_j(x) + \dots$$

is globally negative, where the  $s_i$  and  $r_{ij}$  are sums-of-squares

## **Optimization Problems**

We can combine this with a maximization over t

maximize t subject to  $t-f_0(x)+\sum_{i=1}^m s_i(x)f_i(x)+\sum_{i=1}^m r_{ij}(x)f_i(x)f_j(x)\leq 0 \text{ for all } x$ 

 $s_i, r_{ij}$  are sums-of-squares

- ullet The variables here are (coefficients of) the polynomials  $s_i, r_i$
- We will see later how to approach this kind of problem using semidefinite programming